

CANNA TERRA

GROWSCHEDULE



DAYS AND WEEKS	VEGETATIVE PHASE				GENERATIVE PHASE						
	ROOTING	I	II		I			II	III	IV	
	D1-5	D6-25	W1	W2	W3	W4	W5	W6	W7	W8	W9
LIGHT HOURS	18	18	12	12	12	12	12	12	12	12	12
Dosages in ml per 10 L											
CANNA TERRA VEGA	32 - 49	45 - 62	54 - 71	54 - 71							
CANNA TERRA FLORES					57 - 76	57 - 76	57 - 76	57 - 76	57 - 76	34 - 53	
CANNA RHIZOTONIC	40	20	20	20	5	5	5	5	5		
CANNAZYM		25	25	25	25	25	25	25	25	25 - 50 *	25 - 50
CANNABOOST			20 **	20	20 - 40	20 - 40	20 - 40	20 - 40	20 - 40	20 - 40	20 - 40
CANNA PK 13/14								15			
CANNA CALMAG AGENT***	0 - 7	0 - 7	0 - 7	0 - 7	0 - 7	0 - 7	0 - 7	0 - 7	0 - 7	0 - 7	0 - 7
CANNA FLUSH									40		40
mS/cm											
EC WATER ***	0,4	0,4	0,4	0,4	0,4	0,4	0,4	0,4	0,4	0,4	0,4
EC NUTRITION	0,8 - 1,2	1,1 - 1,5	1,3 - 1,7	1,3 - 1,7	1,3 - 1,7	1,3 - 1,7	1,3 - 1,7	1,7 - 2,1	1,3 - 1,7	0,8 - 1,2	
EC TOTAL	1,2 - 1,6	1,5 - 1,9	1,7 - 2,1	1,7 - 2,1	1,7 - 2,1	1,7 - 2,1	1,7 - 2,1	2,1 - 2,5	1,7 - 2,1	1,2 - 1,6	0,4

This cultivation schedule is intended for plants that develop flowers or fruits, starting from a one-week-old cutting or seedling. The duration of each phase may vary depending on the species.

- * Double CANNAZYM dosage to 50 ml/10 L, if substrate is reused.
- ** Increase CANNABOOST to a maximum of 40 ml/10 L through flowering for extra strength.
- *** CANNA CALMAG AGENT rate depends on the EC_{water}, which ideally should be 0,2-0,4 mS/cm. If the starting water has an EC of 0,0, Calmag could be applied at the maximum rate to achieve an EC of 0,4 mS/cm. If starting water is already higher than 0,0, a lower dose of Calmag should be used. This example schedule assumes an EC_{water} that is (adjusted with Calmag to) 0,4 mS/cm.

For a longer growth phase, repeat the recommended dosages from vegetative Phase I (D6-25). For a longer flowering phase, repeat the recommended dosages from generative Phase I (W3-W5). Add CANNA PK 13/14 in the period 3-4 weeks before harvest. Recommended pH is between 5.8 and 6.2.

